# (19) World Intellectual Property Organization International Bureau





(43) International Publication Date 24 July 2003 (24.07.2003)

**PCT** 

(10) International Publication Number WO 03/060050 A1

- (51) International Patent Classification<sup>7</sup>: 3/50, 3/34, 3/20, 3/18, 1/74
- C11D 1/83,
- (21) International Application Number: PCT/US03/00287
- (22) International Filing Date: 7 January 2003 (07.01.2003)
- (25) Filing Language:

English

(26) Publication Language:

English

- (30) Priority Data: 10/040,667
- 8 January 2002 (08.01.2002) US
- (71) Applicant: COLGATE-PALMOLIVE COMPANY [US/US]; 300 Park Avenue, New York, NY 10022 (US).
- (72) Inventors: ASZMAN, Harry; 43 Baird Road, Englishtown, NJ 07726 (US). KUGLER, Alison; 3 Orchard Parkway, Morganville, NJ 07751 (US).
- (74) Agents: NANFELDT, Richard, E. et al.; Colgate-PalmoliveCompany, 909 River Road, P.O. Box 1343, Piscataway, NJ 08855-1343 (US).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

3/060050 41

(54) Title: ALL PURPOSE LIQUID CLEANING COMPOSITIONS

(57) Abstract: A hard surface liquid detergent with desirable cleansing properties comprises an anionic surfactant, a nonionic surfactant, a solubilizing agent, a solubilizer and water.

### ALL PURPOSE LIQUID CLEANING COMPOSITIONS

#### Field of Invention

10

15

20

25

This invention relates to an all purpose liquid cleaning composition designed in particular for cleaning hard surfaces and which is effective in removing particular and grease soil in leaving unrinsed surfaces with a shiny appearance.

#### Background of the Invention

This invention relates to an improved all-purpose liquid cleaning composition designed in particular for cleaning hard surfaces and which is effective in removing grease soil and/or bath soil and in leaving unrinsed surfaces with a shiny appearance.

In recent years all-purpose liquid detergents have become widely accepted for cleaning hard surfaces, e.g., painted woodwork and panels, tiled walls, wash bowls, bathtubs, linoleum or tile floors, washable wall paper, etc.. Such all-purpose liquids comprise clear and opaque aqueous mixtures of water-soluble synthetic organic detergents and water-soluble detergent builder salts. In order to achieve comparable cleaning efficiency with granular or powdered all-purpose cleaning compositions, use of water-soluble inorganic phosphate builder salts was favored in the prior art all-purpose liquids. For example, such early phosphate-containing compositions are described in U.S. Patent Nos. 2,560,839; 3,234,138; 3,350,319; and British Patent No. 1,223,739.

In view of the environmentalist's efforts to reduce phosphate levels in ground water, improved all-purpose liquids containing reduced concentrations of inorganic phosphate builder salts or non-phosphate builder salts have appeared. A particularly useful self-opacified liquid of the latter type is described in U.S. Patent No. 4,244,840.

However, these prior art all-purpose liquid detergents containing detergent builder salts or other equivalent tend to leave films, spots or streaks on cleaned unrinsed surfaces, particularly shiny surfaces. Thus, such liquids require thorough rinsing of the cleaned surfaces which is a time-consuming chore for the user.

In order to overcome the foregoing disadvantage of the prior art all-purpose liquid, U.S. Patent No. 4,017,409 teaches that a mixture of paraffin sulfonate and a

10

15

20

25

reduced concentration of inorganic phosphate builder salt should be employed. However, such compositions are not completely acceptable from an environmental point of view based upon the phosphate content. On the other hand, another alternative to achieving phosphate-free all-purpose liquids has been to use a major proportion of a mixture of anionic and nonionic detergents with minor amounts of glycol ether solvent and organic amine as shown in U.S. Patent NO. 3,935,130. Again, this approach has not been completely satisfactory and the high levels of organic detergents necessary to achieve cleaning cause foaming which, in turn, leads to the need for thorough rinsing which has been found to be undesirable to today's consumers.

Another approach to formulating hard surfaced or all-purpose liquid detergent composition where product homogeneity and clarity are important considerations involves the formation of oil-in-water (o/w) microemulsions which contain one or more surface-active detergent compounds, a water-immiscible solvent (typically a hydrocarbon solvent), water and a "cosurfactant" compound which provides product stability. By definition, an o/w microemulsion is a spontaneously forming colloidal dispersion of "oil" phase particles having a particle size in the range of 25 to 800 Å in a continuous aqueous phase.

In view of the extremely fine particle size of the dispersed oil phase particles, microemulsions are transparent to light and are clear and usually highly stable against phase separation.

Patent disclosures relating to use of grease-removal solvents in o/w microemulsions include, for example, European Patent Applications EP 0137615 and EP 0137616 - Herbots et al; European Patent Application EP 0160762 - Johnston et al; and U.S. Patent No. 4,561,991 - Herbots et al. Each of these patent disclosures also teaches using at least 5% by weight of grease-removal solvent.

It also is known from British Patent Application GB 2144763A to Herbots et al, published March 13, 1985, that magnesium salts enhance grease-removal performance of organic grease-removal solvents, such as the terpenes, in o/w

15

20

microemulsion liquid detergent compositions. The compositions of this invention described by Herbots et al. require at least 5% of the mixture of grease-removal solvent and magnesium salt and preferably at least 5% of solvent (which may be a mixture of water-immiscible non-polar solvent with a sparingly soluble slightly polar solvent) and at least 0.1% magnesium salt.

However, since the amount of water immiscible and sparingly soluble components which can be present in an o/w microemulsion, with low total active ingredients without impairing the stability of the microemulsion is rather limited (for example, up to 18% by weight of the aqueous phase), the presence of such high quantities of grease-removal solvent tend to reduce the total amount of greasy or oily soils which can be taken up by and into the microemulsion without causing phase separation.

The following representative prior art patents also relate to liquid detergent cleaning compositions in the form of o/w microemulsions: U.S. Patents No. 4,472,291 - Rosario; U.S. Patent No. 4,540,448 - Gauteer et al; U.S. Patent No. 3,723,330 - Sheflin; etc.

Liquid detergent compositions which include terpenes, such as d-limonene, or other grease-removal solvent, although not disclosed to be in the form of o/w microemulsions, are the subject matter of the following representative patent documents: European Patent Application 0080749; British Patent Specification 1,603,047; and U.S. Patent Nos. 4,414,128 and 4,540,505. For example, U.S. Patent No. 4,414,128 broadly discloses an aqueous liquid detergent composition characterized by, by weight:

- (a) from 1% to 20% of a synthetic anionic, nonionic, amphoteric or
   zwitterionic surfactant or mixture thereof;
  - (b) from 0.5% to 10% of a mono- or sesquiterpene or mixture thereof, at a weight ratio of (a):(b) being in the range of 5:1 to 1:3; and
  - (c ) from 0.5% 10% of a polar solvent having a solubility in water at  $15^{\circ}$ C in the range of from 0.2% to 10%. Other ingredients present in the formulations disclosed

15

20

25

in this patent include from 0.05% to 2% by weight of an alkali metal, ammonium or alkanolammonium soap of a C<sub>13</sub>-C<sub>24</sub> fatty acid; a calcium sequestrant from 0.5% to 13% by weight; non-aqueous solvent, e.g., alcohols and glycol ethers, up to 10% by weight; and hydrotropes, e.g., urea, ethanolamines, salts of lower alkylaryl sulfonates, up to 10% by weight. All of the formulations shown in the Examples of this patent include relatively large amounts of detergent builder salts which are detrimental to surface shine.

Furthermore, the present inventors have observed that in formulations containing grease-removal assisting magnesium compounds, the addition of minor amounts of builder salts, such as alkali metal polyphosphates, alkali metal carbonates, nitrilotriacetic acid salts, and so on, tends to make it more difficult to form stable microemulsion systems.

U.S. Patent 5,082,584 discloses a microemulsion composition having an anionic surfactant, a cosurfactant, nonionic surfactant, perfume and water; however, these compositions do not possess the ecotoxicity and the improved interfacial tension properties as exhibited by the compositions of the instant invention.

A number of patents teach esterified ethoxylated glycerol compounds for various applications. These patents are Great Britain 1,453,385; Japan 59-1600 and Japan 58-206693 and European Patent Application 0586,323A1. These publications fail to appreciate that a mixture of esterified ethoxylated glycerol and nonesterified ethoxylated glycerol, when used in a hard surface cleaning composition, functions as a grease release agent.

#### Summary of the Invention

It has now been found that an all purpose hard surface liquid detergent which has desirable cleaning properties can be formulated with an anionic surfactant, a nonionic surfactant and a solubilizer.

An object of this invention is to provide an all purpose hard surface liquid detergent composition comprises a sulfate and/or sulfonate anionic surfactant, a nonionic surfactant, a solubilizing system and water, wherein the composition does not

contain any N-alkyl aldonamide, zwitterionic surfactant, silicas, abrasives, alkali metal carbonates, alkaline earth metal carbonates, alkyl glycine surfactant or a cyclic imidinium surfactant.

Additional objects, advantages and novel features of the invention will be set

forth in part in the description which follows, and in part will become apparent to those skilled in the art upon examination of the following or may be learned by practice of the invention. The objects and advantages of the invention may be realized and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims.

## 10 <u>Detailed Description of the Invention</u>

15

25

The all purpose hard surface liquid compositions of the instant invention comprises approximately by weight:

- (a) 0.1% to 5%, more preferably 0.25% to 3% of an alkali metal or alkaline earth salt of an anionic surfactant;
  - (b) 0.1% to 6%, more preferably 0.25% to 3% of a nonionic surfactant;
- (c) 0.1% to 2% of a solubilizer selected from the group consisting of Polysorbate 80 and PEG30 castor oil and mixtures thereof;
- (d) 0.1% to 4% of perfume, essential oil, a water insoluble organic compound such as an ester or a water insoluble material such as terpene or essential oils; and
- 20 (e) the balance being water.

The instant compositions do not contain an N-alkyl aldonamide, choline chloride or buffering system which is a nitrogerious buffer which is ammonium or alkaline earth carbonate, guanidine derivates, alkoxylalkyl amines and alkyleneamines C<sub>3</sub>-C<sub>7</sub> alkyl and alkenyl monobasic and dibasic acids such as C<sub>4</sub>-C<sub>7</sub> aliphatic carboxylic diacids which do not contain a hydroxy group, phosphoric acid, amino alkylene phosphonic acid and the composition is pourable and is not a gel and the composition has a complex viscosity at 1 rads-1 of less than 0.4 Pascal seconds.

The anionic sulfonate surfactants which may be used in the detergent of this invention are water soluble and include the magnesium, sodium, potassium,

10

15

20

25

ammonium and ethanolammonium salts of linear C8-C16 alkyl benzene sulfonates; C10-C20 paraffin sulfonates, alpha olefin sulfonates containing about 10-24 carbon atoms and C8-C18 ethoxylated alkyl ether sulfates and mixtures thereof. The preferred anionic sulfonate surfactant is a sodium salt of a C12-14 alkyl benzene sulfonate.

The higher alkyl mononuclear aromatic sulfonates, such as the higher alkylbenzene sulfonates containing 9 to 18 or preferably 9 to 16 carbon atoms in the higher alkyl group in a straight or branched chain. A preferred alkylbenzene sulfonate is a linear alkylbenzene sulfonate having a higher content of 3-phenyl (or higher) isomers and a correspondingly lower content (well below 50%) of 2-phenyl (or lower) isomers, such as those sulfonates wherein the benzene ring is attached mostly at the 3 or higher (for example 4, 5, 6 or 7) position of the alkyl group and the content of the isomers in which the benzene ring is attached in the 2 or 1 position is correspondingly low. Preferred materials are set forth in U.S. Patent 3,320,174, especially those in which the alkyls are of 10 to 13 carbon atoms.

Examples of suitable other sulfonated anionic detergents are the well known. The paraffin sulfonates may be monosulfonates or di-sulfonates and usually are mixtures thereof, obtained by sulfonating paraffins of 10 to 20 carbon atoms. Preferred paraffin sulfonates are those of C<sub>12-18</sub> carbon atoms chains, and more preferably they are of C<sub>14-17</sub> chains. Paraffin sulfonates that have the sulfonate group(s) distributed along the paraffin chain are described in U.S. Patents 2,503,280; 2,507,088; 3,260,744; and 3,372,188; and also in German Patent 735,096. Such compounds may be made to specifications and desirably the content of paraffin sulfonates outside the C<sub>14-17</sub> range will be minor and will be minimized, as will be any contents of di- or polysulfonates.

The C8-18 ethoxylated alkyl ether sulfate surfactants have the structure

## R-(OCHCH<sub>2</sub>)<sub>n</sub>OSO<sub>3</sub>M

wherein n is about 1 to about 22 more preferably 1 to 3 and R is an alkyl group having about 8 to about 18 carbon atoms, more preferably 12 to 15 and natural cuts, for

15

20

25

example, C<sub>12-14</sub> or C<sub>12-16</sub> and M is an ammonium cation or a metal cation, most preferably sodium.

The ethoxylated alkyl ether sulfate may be made by sulfating the condensation product of ethylene oxide and C8-10 alkanol, and neutralizing the resultant product. The ethoxylated alkyl ether sulfates differ from one another in the number of carbon atoms in the alcohols and in the number of moles of ethylene oxide reacted with one mole of such alcohol. Preferred ethoxylated alkyl ether polyethenoxy sulfates contain 12 to 15 carbon atoms in the alcohols and in the alkyl groups thereof, e.g., sodium myristyl (3 EO) sulfate.

Ethoxylated C8-18 alkylphenyl ether sulfates containing from 2 to 6 moles of ethylene oxide in the molecule are also suitable for use in the invention compositions. These detergents can be prepared by reacting an alkyl phenol with 2 to 6 moles of ethylene oxide and sulfating and neutralizing the resultant ethoxylated alkylphenol.

The nonionic surfactants used in the instant compositions include the condensation products of a higher alcohol (e.g., an alkanol containing about 8 to 16 carbon atoms in a straight or branched chain configuration) condensed with about 4 to 20 moles of ethylene oxide, for example, lauryl or myristyl alcohol condensed with about 16 moles of ethylene oxide (EO), tridecanol condensed with about 6 to 15 moles of EO, myristyl alcohol condensed with about 10 moles of EO per mole of myristyl alcohol, the condensation product of EO with a cut of coconut fatty alcohol containing a mixture of fatty alcohols with alkyl chains varying from 10 to about 14 carbon atoms in length and wherein the condensate contains either about 6 moles of EO per mole of total alcohol or about 9 moles of EO per mole of alcohol and tallow alcohol ethoxylates containing 6 EO to 11 EO per mole of alcohol.

A preferred group of the foregoing nonionic surfactants are the Neodol ethoxylates (Shell Co.), which are higher aliphatic, primary alcohol containing about 9-15 carbon atoms, such as C9-C11 alkanol condensed with 4 to 10 moles of ethylene oxide (Neodol 91-8 or Neodol 91-5), C12-13 alkanol condensed with 6.5 moles ethylene oxide (Neodol 23-6.5), C12-15 alkanol condensed with 12 moles ethylene

10

20

25

oxide (Neodol 25-12), C14-15 alkanol condensed with 13 moles ethylene oxide (Neodol 45-13), and the like. Such ethoxamers have an HLB (hydrophobic lipophilic balance) value of about 8 to 15 and give good O/W emulsification, whereas ethoxamers with HLB values below 7 contain less than 4 ethyleneoxide groups and tend to be poor emulsifiers and poor detergents.

Additional satisfactory water soluble alcohol ethylene oxide condensates are the condensation products of a secondary aliphatic alcohol containing 8 to 18 carbon atoms in a straight or branched chain configuration condensed with 5 to 30 moles of ethylene oxide. Examples of commercially available nonionic detergents of the foregoing type are C<sub>11</sub>-C<sub>15</sub> secondary alkanol condensed with either 9 EO (Tergitol 15-S-9) or 12 EO (Tergitol 15-S-12) marketed by Union Carbide.

Another water soluble nonionic surfactants which can be utilized in this invention are an aliphatic ethoxylated/propoxylated nonionic surfactants which are depicted by the formula:

15 
$$R = O = (CH_2CH_2O)_{\overline{X}} = (CH_2CH_2CH_2O)_{\overline{Y}} + O$$

wherein R is a branched chain alkyl group having about 10 to about 16 carbon atoms, preferably an isotridecyl group and x and y are independently numbered from 1 to 20. A preferred ethoxylated/propoxylated nonionic surfactant is Plurafac® 300 manufactured by BASF.

As used herein and in the appended claims the term "perfume" is used in its ordinary sense to refer to and include any non-water soluble fragrant substance or mixture of substances including natural (i.e., obtained by extraction of flower, herb, blossom or plant), artificial (i.e., mixture of natural oils or oil constituents) and synthetically produced substance) odoriferous substances. Typically, perfumes are complex mixtures of blends of various organic compounds such as alcohols, aldehydes, ethers, aromatic compounds and varying amounts of essential oils (e.g.,

10

20

terpenes) such as from 0% to 80%, usually from 10% to 70% by weight, the essential oils themselves being volatile odoriferous compounds and also serving to dissolve the other components of the perfume.

In the present invention the precise composition of the perfume is of no particular consequence to cleaning performance so long as it meets the criteria of water immiscibility and having a pleasing odor. Naturally, of course, especially for cleaning compositions intended for use in the home, the perfume, as well as all other ingredients, should be cosmetically acceptable, i.e., non-toxic, hypoallergenic, etc.. The instant compositions show a marked improvement in ecotoxocity as compared to existing commercial products.

In place of the perfume in either the microemulsion composition or the all purpose hard surface cleaning composition at the same previously defined concentrations that the perfume was used in either the microemulsion or the all purpose hard surface cleaning composition one can employ an essential oil or a water insoluble hydrocarbon having 6 to 18 carbon such as a paraffin or isoparaffin.

Suitable essential oils are selected from the group consisting of:

Anethole 20/21 natural, Aniseed oil china star, Aniseed oil globe brand, Balsam (Peru),
Basil oil (India), Black pepper oil, Black pepper oleoresin 40/20, Bois de Rose (Brazil)
FOB, Borneol Flakes (China), Camphor oil, White, Camphor powder synthetic
technical, Cananga oil (Java), Cardamom oil, Cassia oil (China), Cedarwood oil (China)
BP, Cinnamon bark oil, Cinnamon leaf oil, Citronella oil, Clove bud oil, Clove leaf,
Coriander (Russia), Coumarin 69°C (China), Cyclamen Aldehyde, Diphenyl oxide, Ethyl
vanilin, Eucalyptol, Eucalyptus oil, Eucalyptus citriodora, Fennel oil, Geranium oil,
Ginger oil, Ginger oleoresin (India), White grapefruit oil, Guaiacwood oil, Gurjun
balsam, Heliotropin, Isobornyl acetate, Isolongifolene, Juniper berry oil, L-methyl
acetate, Lavender oil, Lemon oil, Lemongrass oil, Lime oil distilled, Litsea Cubeba oil,
Longifolene, Menthol crystals, Methyl cedryl ketone, Methyl chavicol, Methyl salicylate,
Musk ambrette, Musk ketone, Musk xylol, Nutmeg oil, Orange oil, Patchouli oil,
Peppermint oil, Phenyl ethyl alcohol, Pimento berry oil, Pimento leaf oil, Rosalin,

Zestorai<sup>™</sup>.

20

25

Sandalwood oil, Sandenol, Sage oil, Clary sage, Sassafras oil, Spearmint oil, Spike lavender, Tagetes, Tea tree oil, Vanilin, Vetyver oil (Java), Wintergreen, Allocimene, Arbanex<sup>™</sup>, Arbanol<sup>®</sup>, Bergamot oils, Camphene, Alpha-Campholenic aldehyde, I-Carvone, Cineoles, Citral, Citronellol Terpenes, Alpha-Citronellol, Citronellyl Acetate, 5 Citronellyl Nitrile, Para-Cymene, Dihydroanethole, Dihydrocarveol, d-Dihydrocarvone, Dihydrolinalool, Dihydromyrcene, Dihydromyrcenol, Dihydromyrcenyl Acetate, Dihydroterpineol, Dimethyloctanal, Dimethyloctanol, Dimethyloctanyl Acetate, Estragole, Ethyl-2 Methylbutyrate, Fenchol, Ferniol™, Florilys™, Geraniol, Geranyl Acetate, Geranyl Nitrile, Glidmint<sup>™</sup> Mint oils, Glidox<sup>™</sup>, Grapefruit oils, trans-2-Hexenal, 10 trans-2-Hexenol, cis-3-Hexenyl Isovalerate, cis-3-Hexanyl-2-methylbutyrate, Hexyl Isovalerate, Hexyl-2-methylbutyrate, Hydroxycitronellal, Ionone, Isobornyl Methylether, Linalool, Linalool Oxide, Linalyl Acetate, Menthane Hydroperoxide, I-Methyl Acetate, Methyl Hexyl Ether, Methyl-2-methylbutyrate, 2-Methylbutyl Isovalerate, Myrcene, Nerol, Neryl Acetate, 3-Octanol, 3-Octyl Acetate, Phenyl Ethyl-2-methylbutyrate, 15 Petitgrain oil, cis-Pinane, Pinane Hydroperoxide, Pinanol, Pine Ester, Pine Needle oils, Pine oil, alpha-Pinene, beta-Pinene, alpha-Pinene Oxide, Plinol, Plinyl Acetate, Pseudo Ionone, Rhodinol, Rhodinyl Acetate, Spice oils, alpha-Terpinene, gamma-Terpinene, Terpinene-4-OL, Terpineol, Terpinolene, Terpinyl Acetate, Tetrahydrolinalool, Tetrahydrolinalyl Acetate, Tetrahydromyrcenol, Tetralol®, Tomato oils, Vitalizair,

The instant all purpose cleaning composition contains about 0.1 wt. % to about 6 wt. %, of at least one solubilizing agent selected from the group consisting of a C<sub>2-5</sub> mono, dihydroxy or polyhydroxy alkanols such as ethanol, isopropanol, glycerol ethylene glycol, diethylene glycol, propylene glycol and alkali metal cumene or xylene sulfonates such as sodium cumene sulfonate and sodium xylene sulfonate and mixtures thereof. The solubilizing agents are included in order to control low temperature cloud clear properties and aide in cleaning.

The solubilizer agent which is used in the instant compositions are selected from the group consisting of PEG30 castor oil, Polysorbate 80, cocoamido propyl dimethyl

15

20

25

betaine, cocoamido propyl dimethyl amine oxide, Icanol TDA-8, Pluronic F127, Icanol DA-6, PEG40 castor oil and mixtures thereof.

The instant formulas explicitly exclude alkali metal silicates and alkali metal builders such as alkali metal polyphosphates, alkali metal carbonates and alkali metal phosphonates because these materials, if used in the instant composition, would cause the composition to have a high pH as well as leaving residue on the surface being cleaned.

The final essential ingredient in the inventive microemulsion or nonmicroemulsion compositions having improved interfacial tension properties is water. The proportion of water in the compositions generally is in the range of 80% to 98%, preferably 85% to 97% by weight of the cleaning composition.

In addition to the above-described essential ingredients required for the formation of the microemulsion composition, the compositions of this invention may often and preferably do contain one or more additional ingredients which serve to improve overall product performance.

One such ingredient is an inorganic or organic salt of oxide of a multivalent metal cation, particularly Mg<sup>++</sup>. The metal salt or oxide provides several benefits including improved cleaning performance in dilute usage, particularly in soft water areas, and minimized amounts of perfume required to obtain the microemulsion state. Magnesium sulfate, either anhydrous or hydrated (e.g., heptahydrate), is especially preferred as the magnesium salt. Good results also have been obtained with magnesium oxide, magnesium chloride, magnesium acetate, magnesium propionate and magnesium hydroxide. These magnesium salts can be used with formulations at neutral or acidic pH since magnesium hydroxide will not precipitate at these pH levels.

Although magnesium is the preferred multivalent metal from which the salts (inclusive of the oxide and hydroxide) are formed, other polyvalent metal ions also can be used provided that their salts are nontoxic and are soluble in the aqueous phase of the system at the desired pH level.

10

15

20

The compositions can include from 0% to 2.5%, preferably from 0.1% to 2.0% by weight of the composition of a C8-C22 fatty acid or fatty acid soap as a foam suppressant.

The addition of fatty acid or fatty acid soap provides an improvement in the rinseability of the composition whether applied in neat or diluted form. Generally, however, it is necessary to increase the level of cosurfactant to maintain product stability when the fatty acid or soap is present. If more than 2.5 wt. % of a fatty acid is used in the instant compositions, the composition will become unstable at low temperatures as well as having an objectionable smell.

As example of the fatty acids which can be used as such or in the form of soap, mention can be made of distilled coconut oil fatty acids, "mixed vegetable" type fatty acids (e.g. high percent of saturated, mono-and/or polyunsaturated C<sub>18</sub> chains); oleic acid, stearic acid, palmitic acid, eiocosanoic acid, and the like, generally those fatty acids having from 8 to 22 carbon atoms being acceptable.

A proton donating agent such as a hydroxy aliphatic acid can optionally be used in the cleaning composition at a concentration of about 0 wt. % to about 5 wt. %, more preferably about 0.1 wt. % to about 4 wt. %. The hydroxy aliphatic acid used in the instant composition is selected from the group consisting of glycolic acid, salicylic acid, tartaric acid, citric acid and lactic acid and mixtures thereof.

A polyethylene glycol can be optionally used in the instant composition at a concentration of 0 to 6 wt. %, more preferably 0.25 wt. % to 5 wt. % and the polyethylene glycol has a molecular weight of 200 to 1,000, wherein the polyethylene glycol has the structure

HO(CH2CH2O)nH

25 wherein n is 4 to 25.

The liquid cleaning composition of this invention may, if desired, also contain other components either to provide additional effect or to make the product more attractive to the consumer. The following are mentioned by way of example: Colors or dyes in amounts up to 0.5% by weight; preservatives or antioxidizing agents, such as

10

20

formalin, 5-bromo-5-nitro-dioxan-1,3; 5-chloro-2-methyl-4-isothaliazolin-3-one, 2,6-ditert.butyl-p-cresol, etc., in amounts up to 2% by weight; and pH adjusting agents, such as sulfuric acid or sodium hydroxide, as needed. Furthermore, if opaque compositions are desired, up to 4% by weight of an opacifier may be added.

In final form, the instant compositions exhibit stability at reduced and increased temperatures. More specifically, such compositions remain clear and stable in the range of 5°C to 50°C, especially 10°C to 43°C. Such compositions exhibit a pH of 3 to 8.0 which is achieved by the addition of caustic soda or alkanolamide. The liquid compositions are readily pourable and exhibit a viscosity in the range of 6 to 400 milliPascal . second (mPas.) as measured at 25°C. with a Brookfield RVT Viscometer using a #2 spindle rotating at 50 RPM.

The following examples illustrate liquid cleaning compositions of the described invention. Unless otherwise specified, all percentages are by weight. The exemplified compositions are illustrative only and do not limit the scope of the invention. Unless otherwise specified, the proportions in the examples and elsewhere in the specification are by weight.

Example 1

The following compositions in wt. % were prepared by simple mixing procedure:

	Α	В	Standard Formula C
Demineralized water	Bal.	Bal.	Bal.
C10-C13 alkyl linear benzene sulfonate	1.5	1.5	1.6
C9-C11 alcohol EO7.5-8:1	1.0	1.0	
Formalin	0.2	0.2	0.2
EDTA Na4-Dehydrate	0.2	0.3	0.2
Polysorbate 80	0.45		-
PEG30 castor oil		0.24	
Perfume	1.2	1.2	1.2
C9-C11 alcohol 7EO			3.0

В Standard Formula 7.0 7.0 7.0 Appearance clear clear clear Degreasing diluted slightly better slightly better standard Degreasing neat slightly better slightly better standard Residue parity parity standard

#### What Is Claimed:

5

15

- 1. A microemulsion composition comprising:
- (a) 0.1 wt. % to 5 wt. % of a nonionic surfactant;
- (b) 0.1 wt. % to 5 wt. % of an anionic surfactant:
- (c) 0.1 wt. % to 4 wt. % of a water insoluble hydrocarbon, essential oil or a perfume;
  - (d) 0.1 wt. % to 2 wt. % of a solubilizer; and
  - (e) the balance being water.
- 2. The composition of Claim 1 which further contains a salt of a multivalent metal cation in an amount sufficient to provide from 0.5 to 1.5 equivalents of said cation per equivalent of said anionic detergent.
  - The composition of Claim 1 further including a fatty acid has 8 to 22 carbon atoms.
    - 4. The composition of Claim 1 further including polyethylene glycol.
    - 5. The composition of Claim 4 further including a hydroxy aliphatic acid.
  - 6. The composition of Claim 1, wherein said anionic surfactant is a linear alkyl benzene sulfonate.
  - 7. The composition of Claim 1, wherein said solubilizer is selected from the group consisting of PEG30 castor oil and Polysorbate 80.
- 8. The composition of Claim 7, wherein said solubilizing agent is selected from the group consisting of an alkanol, alkali metal xylene sulfonate and alkali metal cumene sulfonate and mixtures thereof.

Internation No PCT/US 03/00287

A CLASS	IFICATION OF SUBJECT MATTER				
ÎPC 7	C11D1/83 C11D3/50 C11D3/ C11D1/74	<b>′</b> 34 ·	C11D3/20	C110	03/18
According	io international Patent Classification (IPC) or to both national class	sification a	nd IPC		
	SEARCHED				
IPC /	ocumentation searched (classification system followed by classifi ${\tt C11D}$				
	ation searched other than minimum documentation to the extent th				
	data base consulted during the International search (name of data ternal, WPI Data, PAJ	base and	, where practical, s	earch terms use	<b>d</b> )
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the	relevant p	assages		Relevant to claim No.
X .	WO 98 00497 A (COLGATE PALMOLIV 8 January 1998 (1998-01-08) claims; examples 1,6,7	E CO)			1-8
X .	US 6 291 418 B1 (MONDIN MYRIAM 18 September 2001 (2001-09-18) claims; example 1	ET A	-)	.	1-8
X	EP 0 316 726 A (COLGATE PALMOLI 24 May 1989 (1989-05-24) claims; examples 1,6,710	VE CO	*		1-8
X	EP 0 672 747 A (COLGATE PALMOLI 20 September 1995 (1995-09-20) claims; example 1	/E CO)			1-8
	· .	-/			
X Furth	er documents are listed in the continuation of box C.	X	Patent family men	mbers are listed	In annex.
Special call	egories of cited documents:		<del></del>		
'A' documer	nt defining the general state of the art which is not ared to be of particular relevance	di	priority date and no	it in conflict with	mational filing date the application but sory underlying the
"E" earlier de filing da	ocument but published on or after the international site	'X' doc	ument of particular	relevance; the cl	lalmed Invention
"L" document which is	it which may throw doubts on priority claim(s) or s clied to establish the publication date of another	inv	olve an inventive si	novel or cannot sep when the doc	be considered to current is taken alone
CHERON	or other special reason (as specified) ni referring to an oral disclosure, use, exhibition or	ca	ument of particular nnot be considered	to involve an inv	BILIVA SIED When the
oiner m	eans	ine on	ints, such combined	with one or mo	re other such docu- is to a person skilled
later the	nt published prior to the international filling date but an the priority date claimed		he art. ument member of th	ne same patent f	amily
Date of the a	ctual completion of the international search		e of malling of the I		
	May 2003		16/05/200	3	
Name and ma	alling address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Aut	horized officer		
	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax. (+31-70) 340-3016		Pentek F		

Internation Application No
PCT/US 03/00287

		PCT/US 03/00287
C.(Continue Calegory •	ation) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the relevant passages	I Sala and a Ma
Jalegory	Cualion of document, with aducation, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 668 346 A (COLGATE PALMOLIVE CO) 23 August 1995 (1995-08-23) claims; examples I,III	1-8
Κ	US 5 776 880 A (MONDIN MYRIAM ET AL) 7 July 1998 (1998-07-07) claims; example 1	1-8
	US 5 415 813 A (YIANAKOPOULOS GEORGES ET AL) 16 May 1995 (1995-05-16) claims; example 1	1-8
	US 5 854 193 A (MONDIN MYRIAM ET AL) 29 December 1998 (1998-12-29) claims; example 1	1-8
	US 6 268 330 B1 (TOUSSAINT CHRISTINE ET AL) 31 July 2001 (2001-07-31) claim 1	. 5
	US 5 447 644 A (GUENIN ERIC P ET AL) 5 September 1995 (1995-09-05) column 9, line 15 -column 10, line 2	7
	•	
	•	
	·	

ai Application No PCT/US 03/00287

				- 1	PC1/US	03/00287
Ci	Patent document ted in search report		Publication date		Patent family member(s)	Publication date
W	0 9800497	A	08-01-1998	AU WO US US	3233897 A 9800497 A2 5985814 A 5942482 A	21-01-1998 08-01-1998 16-11-1999 24-08-1999
U	S 6291418	B1	18-09-2001	US	6136774 A	24-10-2000
			10 03 1001	US	6057279 A	02-05-2000
				US	6004919 A	21-12-1999
				US US	6136773 A 6288019 B1	24-10-2000
				US	6150319 A	11-09-2001 21-11-2000
	• .			AU	1324900 A	05-06-2000
				EP WO	1129173 A2 0029528 A2	05-09-2001 25-05-2000
				ÜS	5994287 A	30-11-1999
EF	P 0316726	Α.	24-05-1989	US	5076954 A	31-12-1991
				AU Br	2506288 A 8805932 A	18-05-1989 01-08-1989
				CA	1337585 A1	21-11-1995
				DK Ep	628488 A	13-05-1989
	•			ΪĹ	0316726 A2 88339 A	24-05-1989 06-09-1992
				MX	169813 B	27-07-1993
				NO NZ	885051 A 226927 A	16-05-1989 25-10-1991
				PT	88977 A ,B	30-11-1989
				US	5108643 A	28-04-1992
· EP	0672747	Α -	20-09-1995	AT AU	201899 T 1355995 A	15-06-2001
				BR	9501075 A	21-09-1995 04-06-1996
				CA DE	2144367 A1	15-09-1995
				DE	69521162 D1 69521162 T2	12-07-2001 28-02-2002
			=	DK	672747 T3	17-09-2001
				EP ES	0672747 A2 2159307 T3	20-09-1995 01-10-2001
				GR	3036552 T3	31-12-2001
				HU NZ	70070 A2 270633 A	28-09-1995
				PT	672747 T	24-06-1997 30-10-2001
				US	5763386 A	09-06-1998
EP	0668346	A	23-08-1995	US AU	5571459 A 680076 B2	05-11-1996
	•			AU	1148995 A	17-07-1997 17-08-1995
				BR ·	9500451 A	26-09-1995
				CA EP	2141926 A1 0668346 A1	08-08-1995 23-08-1995
				HU	70071 A2	28-09-1995
				NZ US	270433 A	25-06-1996
				PL	5561106 A 307113 A1	01-10-1996 10-06-1996
US	5776880	A	07-07-1998	US	5549840 A	27-08-1996
				US	5952281 A	14-09-1999
				AT	207110 T	15-11-2001

PCT/US 03/00287

	·			101/03	03/00287
Patent docume cited in search re		Publication date		Patent family member(s)	Publication date
US 5776880	Α		AU	696196 B2	03-09-1998
			AU	4105996 A	06-06-1996
			BR	9509682 A	30-09-1997
•			CA	2205404 A1	23-05-1996
		•	CN	1170433 A	14-01-1998
			CZ	9701502 A3	14-10-1998
	•		DE	69523330 D1	22-11-2001
		•	DE	69523330 T2	11-07-2002
			DK	791049 T3	11-07-2002
			EP	0791049 A1	27-08-1997
			ΕP	0994180 A1	19-04-2000
			ES	2165926 T3	01-04-2002
			HU	77481 A2	28-05-1998
			PL	320224 A1	15-09-1997
			PT	791049 T	29-04-2002
			RO -	115540 B1	30-03-2000
		•	RU	2147312 C1	10-04-2000
			WO	9615217 A1	
			US	5759983 A	23-05-1996
•			US	5741760 A	02-06-1998
	•		US ·		21-04-1998
			ZA	5861367 A	19-01-1999
			AT	. 9509530 A	09-05-1997
			AU	211757 T 691499 B2	15-01-2002
			AU	6878094 A	21-05-1998
			BR		16-02-1995
			CA	9403153 A 2129399 A1	11-04-1995
			DE	69429600 D1	05-02-1995 14-02-2002
			DK	637629 T3	29-04-2002
			EP	0637629 A1	08-02-1995
			GR	1003582 B2	22-05-2001
			ΗÜ	67925 A2	20-03-1995
			NO	942886 A	06-02-1995
		•	NZ	264113 A	25-06-1996
			PL	304558 A1	06-02-1995
			PL	184267 B1	30-09-2002
			PT	101556 A ,B	04-05-1995
4			RO	114905 B1	30-08-1999
			üs	5985814 A	16-11-1999
•			US	5763386 A	09-06-1998
			US	5731281 A	24-03-1998
			US	5942482 A	24-08-1999
			US	6020296 A	01-02-2000
			US	6017868 A	25-01-2000
			US	5599785 A	04-02-1997
			US	6191090 B1	20-02-2001
US 5415813	A	16-05-1995	AT	181356 T	15-07-1999
•	•		AU	685204 B2	15-01-1998
		*	AU	1178795 A	13-06-1995
			BR	9408120 A	05-08-1997
			DE	69419173 D1	22-07-1999
			EP	0730634 A1	11-09-1996
			NZ	276935 A	26-08-1998
			PL	319156 A1	21-07-1997
			WO	9514763 A1	01-06-1995
•			US	5552089 A	03-09-1996

Interna Application No
PCT/US 03/00287

Patent docu cited in searc	ıment h report	Publication date		Patent family member(s)	Publication date
US 58541	93 A	29-12-1998	US		
	A	F31F1330	AU	5716925 A 727094 B2	10-02-1998
	•				30-11-2000
			AU	9019498 A	08-03-1999
			CA	2300809 A1	25-02-1999
			EP	1005518 A1	07-06-2000
			HU	0004257 A2	28-04-2001
			NZ	502911 A	29-06-2001
			PL	338703 A1	20-11-2000
		•	WO	9909121 A1	25-02-1999
			US	6017868 A	25-01-2000
			US	6191090 B1	
			US	6020296 A	20-02-2001
			AT		01-02-2000
				200793 T	15-05-2001
			AU	680087 B2	17-07-1997
	•		AU	1289195 A	13-06-1995
			BR	9408122 A	05-08-1997
		•	DE	69427154 D1	31-05-2001
			DE	69427154 T2	29-11-2001
•			DK	730636 T3	13-08-2001
			EP	0730636 A1	11-09-1996
			ES	2158072 T3	01-09-2001
			GR	3036281 T3	31-10-2001
			NZ	277602 A	22-09-1997
			PL	314622 A1	
			PT	730636 T	16-09-1996
			WO		30-10-2001
,				9514765 A1	01-06-1995
			AT	211757 T	15-01-2002
			AU	691499 B2	21-05-1998
		•	AU	6878094 A	16-02-1995
			BR	9403153 A	11-04-1995
			CA	2129399 A1	05-02-1995
			DE	69429600 D1	14-02-2002
			DK	637629 T3	29-04-2002
			EP	0637629 A1	08-02-1995
			GR	1003582 B2	22-05-2001
			HU	67925 A2	20-03-1995
			NO	942886 A	06-02-1995
			NZ	264113 A	25-06-1996
	1		PL	304558 A1	
•	•		PT		06-02-1995
			RO	101556 A ,B	04-05-1995
		•		114905 B1	30-08-1999
			US	5952281 A	14-09-1999
			US	5985814 A	16-11-1999
			US	5599785 A	04-02-1997
	•		US	5549840 A	27-08-1996
			US.	5610130 A	11-03-1997
			US	5763386 A	09-06-1998
			US	5759983 A	02-06-1998
		•	US	5741760 A	21-04-1998
US 626833	) B1	31-07-2001	AU	5150900 A	12-12-2000
		<del>-</del>	EP	1180133 Å1	20-02-2002
			WO	0071667 A1	
			US		30-11-2000
				6255269 B1	03-07-2001
			US	6251844 B1	26-06-2001
			US US	6262003 B1 2001009893 A1	17-07-2001
					26-07-2001

Internation Application No
PCT/US 03/00287

Patent document dted in search report	Publication date	Patent family member(s)	Publication date
US 5447644	05-09-1995	CA 2148675 EP 0682104	

# THIS PAGE BLANK (USPTO)